

Patient age, sex and inflammatory bowel disease phenotype associate with course of Primary Sclerosing Cholangitis

International PSC Study Group (IPSCSG); Weismüller, Tobias J; Trivedi, Palak; Bergquist, Annika; Imam, Mohamad H.; Lenzen, Henrike; Ponsioen, Cyriel Y.; Holm, Kristian; Gotthardt, Daniel Nils; Färkkilä, Martti; Marschall, Hanns-ulrich; Thorburn, Douglas; Weersma, Rinse K; Fevery, J; Mueller, T; Chazouillères, Olivier; Schulze, Kornelius; Lazaridis, Konstantinos N; Almer, Sven; Pereira, SP

DOI:

[10.1053/j.gastro.2017.02.038](https://doi.org/10.1053/j.gastro.2017.02.038)

License:

Creative Commons: Attribution-NonCommercial-NoDerivs (CC BY-NC-ND)

Document Version

Peer reviewed version

Citation for published version (Harvard):

International PSC Study Group (IPSCSG), Weismüller, TJ, Trivedi, P, Bergquist, A, Imam, MH, Lenzen, H, Ponsioen, CY, Holm, K, Gotthardt, DN, Färkkilä, M, Marschall, H, Thorburn, D, Weersma, RK, Fevery, J, Mueller, T, Chazouillères, O, Schulze, K, Lazaridis, KN, Almer, S, Pereira, SP, Levy, C, Mason, A, Næss, S, Bowlus, CL, Floreani, A, Halilbasic, E, Yimam, KK, Milkiewicz, P, Beuers, U, Huynh, D, Pares, A, Manser, CN, Dalekos, GN, Eksteen, B, Invernizzi, P, Berg, CP, Kirchner, GI, Sarrazin, C, Zimmer, V, Fabris, L, Braun, F, Marziani, M, Juran, BD, Said, K, Rupp, C, Jokelainen, K, de Valle, M, Saffioti, F, Cheung, A, Trauner, M, Schramm, C, Chapman, RW, Karlsen, TH, Schrupf, E, Strassburg, CP, Manns, MP, Lindor, KD, Hirschfield, G, Hansen, BE & Boberg, KM 2017, 'Patient age, sex and inflammatory bowel disease phenotype associate with course of Primary Sclerosing Cholangitis', *Gastroenterology*, vol. 152, no. 8, pp. 1975-1984.e8. <https://doi.org/10.1053/j.gastro.2017.02.038>

[Link to publication on Research at Birmingham portal](#)

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

Download date: 05. May. 2023

Gastroenterology

Patient Age, Sex, and Inflammatory Bowel Disease Phenotype Associate With Course of Primary Sclerosing Cholangitis

--Manuscript Draft--

Manuscript Number:	GASTRO-D-16-00774R1
Full Title:	Patient Age, Sex, and Inflammatory Bowel Disease Phenotype Associate With Course of Primary Sclerosing Cholangitis
Article Type:	Clinical - Biliary
Section/Category:	Clinical
Corresponding Author:	Tobias Johannes Weismüller University of Bonn Bonn, GERMANY
Corresponding Author Secondary Information:	
Corresponding Author's Institution:	University of Bonn
Corresponding Author's Secondary Institution:	
First Author:	Tobias Johannes Weismüller
First Author Secondary Information:	
Order of Authors:	Tobias Johannes Weismüller
	Palak J. Trivedi
	Annika Bergquist
	Mohamad Imam
	Henrike Lenzen
	Cyriel Y Ponsioen
	Kristian Holm
	Daniel Gotthardt
	Martti A Färkkilä
	Hanns-Ulrich Marschall
	Douglas Thorburn
	Rinse K Weersma
	Johan Fevery
	Tobias Mueller
	Olivier Chazouillères
	Kornelius Schulze
	Konstantinos N Lazaridis
	Sven Almer
	Stephen P Pereira
	Cynthia Levy
	Andrew Mason
	Sigrid Naess

	Christopher L Bowlus
	Annarosa Floreani
	Emina Halilbasic
	Kidist K Yimam
	Piotr Milkiewicz
	Ulrich Beuers
	Dep K Huynh
	Albert Pares
	Christine N Manser
	George N Dalekos
	Bertus Eksteen
	Pietro Invernizzi
	Christoph P Berg
	Gabi I Kirchner
	Christoph Sarrazin
	Vincent Zimmer
	Luca Fabris
	Felix Braun
	Marco Marzioni
	Brian D Juran
	Karouk Said
	Christian Rupp
	Kalle Jokelainen
	Maria Benito de Valle
	Francesca Saffioti
	Angela Cheung
	Michael Trauner
	Christoph Schramm
	Roger W Chapman
	Tom H Karlsen
	Erik Schrumpf
	Christian P Strassburg
	Michael P Manns
	Keith D Lindor
	Gideon M Hirschfield
	Bettina E Hansen
	Kirsten M Boberg
Order of Authors Secondary Information:	
Abstract:	Background & Aims: Primary sclerosing cholangitis (PSC) is an orphan hepatobiliary disorder associated with inflammatory bowel disease (IBD). We aimed to estimate the

risk of disease progression based on distinct clinical phenotypes in a large, international cohort of patients with PSC.

Methods: We performed a retrospective outcome analysis of patients diagnosed with PSC from 1980 through 2010 at 37 centers in Europe, North America, and Australia. For each patient, we collected data on sex, clinician-reported age at and date of PSC and IBD diagnoses, phenotypes of IBD and PSC, and date and indication of IBD-related surgeries. The primary and secondary endpoints were liver transplantation or death (LTD) and hepatopancreatobiliary malignancy, respectively. Cox proportional hazards models were applied to determine the effects of individual covariates on rates of clinical events, with time-to-event analysis ascertained through Kaplan-Meier estimates.

Results: Of the 7121 patients in the cohort, 2616 met the primary endpoint (median time-to-event of 14.5 years) and 721 developed hepatopancreatobiliary malignancy. The most common malignancy was cholangiocarcinoma (n=594); patients of advanced age at diagnosis had an increased incidence, compared with younger patients (incidence rate [IR]: 1.2 per 100 patient-years for patients younger than 20 years old, 6.0 per 100 patient-years for patients 21-30 years old, 9.0 per 100 patient-years for patients 31-40 years old, 14.0 per 100 patient-years for patients 41-50 years old, 15.2 per 100 patient-years for patients 51-60 years old, and 21.0 per 100 patient-years for patients older than 60 years). Of all patients with PSC studied, 65.5% were men, 89.8% had classical or large-duct disease, and 70.0% developed IBD at some point. Assessing the development of IBD as a time-dependent covariate, Crohn's disease (CD) and no IBD (both vs ulcerative colitis [UC]) were associated with a lower risk of LTD (unadjusted hazard ratio [HR], 0.62; $P<.001$ and HR, 0.90; $P=.03$; respectively) and malignancy (HR, 0.68; $P=.008$ and HR, 0.77; $P=.004$, respectively). Small-duct PSC was associated with a lower risk of LTD or malignancy compared with classic PSC (HR, 0.30 and HR, 0.15, respectively; both $P<.001$). Female sex was also associated with a lower risk of LTD or malignancy (HR, 0.88; $P=.002$ and HR, 0.68; $P<.001$, respectively). In multivariable analyses assessing the primary endpoint, small-duct PSC characterized a low-risk phenotype in both sexes (adjusted HR for men, 0.23; $P<.001$ and adjusted HR for women, 0.48; $P=.003$). Conversely, patients with UC had an increased risk of liver disease progression compared to patients with CD (HR, 1.56; $P<.001$) or no IBD (HR, 1.15; $P=.002$).

Conclusions: In an analysis of data from individual patients with PSC worldwide, we found significant variation in clinical course associated with age at diagnosis, sex, and ductal- and IBD subtypes. The survival estimates provided might be used to estimate risk levels for patients with PSC and select patients for clinical trials.

KEY WORDS: risk stratification, immune-mediated liver disease, autoimmune liver disease, cholestasis

Patient Age, Sex, and Inflammatory Bowel Disease Phenotype Associate With Course of Primary Sclerosing Cholangitis

Tobias J. Weismüller,^{1,6} * Palak J. Trivedi,^{2,3} * Annika Bergquist,⁴ Mohamad Imam,^{5,45} Henrike Lenzen,⁶ Cyriel Y. Ponsioen,⁷ Kristian Holm,⁸ Daniel Gotthardt,⁹ Martti A. Färkkilä,¹⁰ Hanns-Ulrich Marschall,¹¹ Douglas Thorburn,¹² Rinse K. Weersma,¹³ Johan Fevery,¹⁴ Tobias Mueller,¹⁵ Olivier Chaouillères,¹⁶ Kornelius Schulze,¹⁷ Konstantinos N. Lazaridis,⁵ Sven Almer,¹⁸ Stephen P. Pereira,¹⁹ Cynthia Levy,²⁰ Andrew Mason,²¹ Sigrid Naess,^{8,47} Christopher L. Bowlus,²² Annarosa Floreani,²³ Emina Halilbasic,²⁴ Kidist K. Yimam,²⁵ Piotr Milkiewicz,^{26,27} Ulrich Beuers,⁷ Dep K. Huynh,²⁸ Albert Pares,²⁹ Christine N. Manser,³⁰ George N. Dalekos,³¹ Bertus Eksteen,³² Pietro Invernizzi,³³ Christoph P. Berg,³⁴ Gabi I. Kirchner,³⁵ Christoph Sarrazin,³⁶ Vincent Zimmer,³⁷ Luca Fabris,³⁸ Felix Braun,³⁹ Marco Marzioni,⁴⁰ Brian D. Juran,⁵ Karouk Said,⁴ Christian Rupp,⁹ Kalle Jokelainen,¹⁰ Maria Benito de Valle,¹¹ Francesca Saffioti,¹² **Angela Cheung,**⁵ Michael Trauner,²⁴ Christoph Schramm,^{17, 41} Roger W. Chapman,^{42,43} Tom H. Karlsen,^{8, 47} Erik Schrumpf,^{8, 47} Christian P. Strassburg,¹ Michael P. Manns,⁶ Keith D. Lindor,^{5,44,46} Gideon M Hirschfield,² Bettina E. Hansen,^{48,49,50*} and Kirsten M. Boberg.^{8, 47*} on behalf of the International PSC Study Group

* These authors have contributed equally to this manuscript.

Affiliations

1. Department of Internal Medicine I, University of Bonn, Germany.
2. National Institute for Health Research (NIHR) Birmingham, Liver Biomedical Research Unit (BRU), University of Birmingham, United Kingdom.
3. Liver Unit, University Hospitals Birmingham Queen Elizabeth, United Kingdom.
4. Center for Digestive Diseases, Division of Hepatology, Karolinska University Hospital and Karolinska Institutet, Stockholm, Sweden.
5. Division of Gastroenterology and Hepatology, Mayo Clinic, Rochester, MN, United States of America.
6. Department of Gastroenterology, Hepatology and Endocrinology, Hannover Medical School, Hannover, Germany.
7. Department of Gastroenterology & Hepatology, Academic Medical Center, Amsterdam, Netherlands.
8. Norwegian PSC Research Center and Section for Gastroenterology, Department of Transplantation Medicine, Division of Surgery, Inflammatory Diseases and Transplantation, Oslo University Hospital, Rikshospitalet, Oslo, Norway.
9. Dept. of Gastroenterology, Infectious Diseases and Intoxications, University Hospital Heidelberg, Heidelberg, Germany.
10. Helsinki University, Clinic of Gastroenterology, Helsinki University Hospital, Helsinki, Finland.
11. Department of Molecular and Clinical Medicine, Sahlgrenska Academy, University of Gothenburg, Gothenburg, Sweden.
12. The Sheila Sherlock Liver Centre and UCL Institute for Liver and Digestive Health, Royal Free Hospital, London, United Kingdom.

13. Department of Gastroenterology and Hepatology, University of Groningen and University Medical Center, Groningen, Netherlands.
14. Department of Hepatology, University Hospital Gasthuisberg, Leuven, Belgium.
15. Department of Internal Medicine, Hepatology and Gastroenterology, Charité Universitätsmedizin Berlin, Campus Virchow Klinikum, Berlin, Germany.
16. Service d'Hépatologie, Hôpital Saint Antoine, Assistance Publique-Hôpitaux de Paris, Faculté de Médecine Pierre et Marie Curie, Paris, France.
17. 1st Department of Medicine, University Medical Center Hamburg Eppendorf, Hamburg, Germany.
18. Division of Gastroenterology and Hepatology, Linköping University, Linköping, Sweden. Current affiliation: Department of Medicine, Karolinska Institutet, Solna, and Center for Digestive Diseases, Karolinska University Hospital, Stockholm, Sweden.
19. Institute for Liver and Digestive Health, UCL, London, United Kingdom.
20. **Division of Hepatology**, University of Miami, Miami, FL, United States.
21. Division of Gastroenterology and Hepatology, University of Alberta, Edmonton, AB, Canada.
22. Division of Gastroenterology and Hepatology, University of California Davis, Davis, CA, United States.
23. Department of Surgery, Oncology and Gastroenterology, University of Padova, Padova, Italy.
24. Division of Gastroenterology and Hepatology, Department of Internal Medicine III, Medical University of Vienna, Austria.
25. Department of Hepatology and Liver Transplantation, California Pacific Medical Center, San Francisco, CA, United States.
26. Department of Clinical and Molecular Biochemistry, Pomeranian Medical University, Szczecin, Poland.

27. Liver and Internal Medicine Unit, Department of General, Transplant and Liver Surgery, Medical University of Warsaw, Poland.
28. Department of Gastroenterology and Hepatology, Royal Adelaide Hospital, Adelaide, SA, Australia.
29. Liver Unit, Hospital Clinic, IDIBAPS, CIBERehd, University of Barcelona, Spain.
30. Division for Gastroenterology and Hepatology, University Hospital Zurich (USZ), Zurich, Switzerland.
31. Department of Medicine and Research Laboratory of Internal Medicine, School of Medicine, University of Thessaly, Larissa, Greece.
32. University of Calgary, Snyder Institute for Chronic Diseases, Alberta, AB, Canada.
33. **Program for Autoimmune Liver Diseases**, International Center for Digestive Health, Department of Medicine and Surgery, University of Milan-Bicocca, Milan, Italy.
34. Department of Gastroenterology, Hepatology, and Infectiology, Medical Clinic, University of Tübingen, Germany.
35. Department of Internal Medicine 1, University Hospital of Regensburg, Regensburg, Germany.
36. Department of Internal Medicine 1, Johann Wolfgang Goethe-University Hospital, Frankfurt, Germany.
37. Department of Medicine II, Saarland University Medical Center, Homburg, Germany.
38. Department of Molecular Medicine, University of Padua School of Medicine, Padua, Italy.
39. Department of General, Visceral, Thoracic, Transplantation and Pediatric Surgery, Campus Kiel, UKSH, Kiel, Germany.
40. Clinic of Gastroenterology, Università Politecnica delle Marche, Ancona, Italy.
41. Martin Zeitz Center for Rare Diseases, University Medical Center Hamburg Eppendorf, Hamburg, Germany.

42. Nuffield Department of Clinical Medicine, University of Oxford, UK
43. Translational Gastroenterology Unit, John Radcliffe Hospital, Oxford, UK
44. Division of Gastroenterology and Hepatology, Mayo Clinic, Phoenix, AZ, USA
45. Department of Internal Medicine, University of North Dakota, USA.
46. Arizona State University, College of Health Solutions, Phoenix, AZ, USA.
47. Institute of Clinical Medicine, University of Oslo, Oslo, Norway
48. Department of Gastroenterology and Hepatology, Erasmus University Medical Center, Rotterdam, The Netherlands.
49. Institute of Health Policy, Management and Evaluation, University of Toronto
50. Toronto Centre for Liver Disease, Toronto General Hospital, Toronto, Canada

Correspondence to:

Dr. T.J. Weismüller (tobias.weismueller@gmx.de)

Or

Prof. K. M. Boberg (kboberg@ous-hf.no)

ABSTRACT

Background & Aims: Primary sclerosing cholangitis (PSC) is an orphan hepatobiliary disorder associated with inflammatory bowel disease (IBD). We aimed to estimate the risk of disease progression based on distinct clinical phenotypes in a large, international cohort of patients with PSC.

Methods: We performed a retrospective outcome analysis of patients diagnosed with PSC from 1980 through 2010 at 37 centers in Europe, North America, and Australia. For each patient, we collected data on sex, clinician-reported age at and date of PSC and IBD diagnoses, phenotypes of IBD and PSC, and date and indication of IBD-related surgeries. The primary and secondary endpoints were liver transplantation or death (LTD) and hepatopancreatobiliary malignancy, respectively. Cox proportional hazards models were applied to determine the effects of individual covariates on rates of clinical events, with time-to-event analysis ascertained through Kaplan-Meier estimates.

Results: Of the 7121 patients in the cohort, 2616 met the primary endpoint (median time-to-event of 14.5 years) and 721 developed hepatopancreatobiliary malignancy. The most common malignancy was cholangiocarcinoma (n=594); patients of advanced age at diagnosis had an increased incidence, compared with younger patients (incidence rate [IR]: 1.2 per 100 patient-years for patients younger than 20 years old, 6.0 per 100 patient-years for patients 21–30 years old, 9.0 per 100 patient-years for patients 31–40 years old, 14.0 per 100 patient-years for patients 41–50 years old, 15.2 per 100 patient-years for patients 51–60 years old, and 21.0 per 100 patient-years for patients older than 60 years). Of all patients with PSC studied, 65.5% were men, 89.8% had classical or large-duct disease, and 70.0% developed IBD at some point. Assessing the development of IBD as a time-dependent covariate, Crohn's disease (CD) and no IBD (both vs ulcerative colitis [UC]) were associated with a lower risk of LTD (unadjusted hazard ratio [HR], 0.62; $P<.001$ and HR, 0.90; $P=.03$; respectively) and malignancy (HR, 0.68; $P=.008$ and HR, 0.77; $P=.004$, respectively). Small-duct PSC was associated with a lower risk of LTD or malignancy compared with classic PSC (HR, 0.30 and HR, 0.15, respectively; both $P<.001$). Female sex was also associated with a lower risk of LTD or malignancy (HR, 0.88; $P=.002$ and HR, 0.68; $P<.001$, respectively). In multivariable analyses assessing the primary endpoint, small-duct PSC characterized a low-risk phenotype in both sexes (adjusted HR for

men, 0.23; $P<.001$ and adjusted HR for women, 0.48; $P=.003$). Conversely, patients with UC had an increased risk of liver disease progression compared to patients with CD (HR, 1.56; $P<.001$) or no IBD (HR, 1.15; $P=.002$).

Conclusions: In an analysis of data from individual patients with PSC worldwide, we found significant variation in clinical course associated with age at diagnosis, sex, and ductal- and IBD subtypes. The survival estimates provided might be used to estimate risk levels for patients with PSC and select patients for clinical trials.

KEY WORDS: risk stratification, immune-mediated liver disease, autoimmune liver disease, cholestasis

INTRODUCTION

Primary sclerosing cholangitis (PSC) is a chronic immune-mediated liver disorder strongly associated with inflammatory bowel disease (IBD).¹ Although rare, PSC carries an ongoing and disproportionate clinical need, with clinical outcomes being determined by the development of end-stage biliary cirrhosis and an independent risk of hepatopancreatobiliary (HPB) malignancy. To date, medical therapies have not been effective,⁸ and liver transplantation remains the only proven life-extending intervention, with 10 – 15% of all transplant activity in Europe now being performed for PSC.^{5–7}

Accurately reporting the natural history of disease remains a critical challenge not only for clinicians, but also industry and regulatory agencies who collectively recognise the need for new therapies and equally appreciate the risks and obstacles in demonstrating patient-benefit against the background of an orphan disease with a relatively variable, often slow clinical course.⁹ Moreover, patients seek reassurance and guidance as to their own prognosis, whereas clinicians wish to confidently recognize those at highest risk of poor outcomes as equally as they strive to reassure individuals with a more favorable prognosis.

To expand upon single-center and single-country descriptors, the International PSC Study Group (IPSCSG) sponsored a multi-center outcome study to model the natural history of the disease. Our primary aim was to evaluate and report the clinical course from a large internationally representative PSC cohort; which included 7,121 patients seen at 37 centres across 17 countries, and encompassing >30-years of clinical observation, 1,696 liver transplants, 920 deaths and 721 incidents of HPB malignancy. In so doing we not only validate the presence of key phenotypic descriptors, but also determine the extent of their interaction and how they may impact the clinical course that patients may experience.

PATIENTS AND METHODS

Study setting and design

We collected and analysed data from well-characterised patients diagnosed with PSC between January 1st 1980 and December 31st 2010, having previously attended or under current clinical follow-up until study completion (June 30th 2014). Any individual with an established diagnosis of PSC (including small-duct disease; sdPSC) in accordance with European or American recommendations^{10–12} was considered eligible for inclusion. When biochemical, serological, and/or histological features of autoimmune hepatitis (AIH) were evident concurrently or sequentially,¹³ the diagnosis of a PSC phenotype with AIH features (PSC/AIH variant) was made according to discretion of the participating center. IBD phenotypes were determined according to local expertise,^{14–16} and classified as ulcerative colitis (UC), Crohn's disease (CD), or indeterminate colitis (IC), in keeping with consensus guidelines.^{17,18}

Data collection

Identification of study participants was performed at a local level, either through a pre-existing and prospectively collected local PSC database; or in a retrospective manner via review of medical records by a named site investigator at a given institution. All individual center data was captured onto a multi-parametric standardised case record form formulated by the IPSCSG, and upon study completion amalgamated into a common 'master' database for downstream analysis. Individual clinical characteristics pertained to patient sex, clinician-reported age at and date of diagnosis of PSC, sub-phenotype and IBD phenotype, date and indication of IBD-related surgical resections, date of LT, date of death and date and type of first HPB malignancy. Patients with sclerosing cholangitis suspected due to alternate aetiologies (e.g. IgG4-related disease, acquired immunodeficiency syndromes, confirmed biliary transporter defects) were excluded from the analysis, as were those with inadequate/unknown

follow-up duration. Upon completion of data capture, all **patient datasets** were checked for plausibility and validity, and duplicated patient entries were removed prior to analysis.

Data interpretation and analysis

All patients were identified at time of diagnosis or during subsequent follow-up. ‘Time zero’ was set from point of diagnosis of first PSC phenotype, with the primary endpoint being **the incidence rate (and associated risk) of LT**, or death (LTD) in non-transplanted patients. Any individual not experiencing a clinical event in this regard was censored at date of last known follow-up. A secondary endpoint of HPB malignancy was also studied, and in this instance the date of first liver transplantation/death, or last date of ‘event-free’ follow-up comprised our censor points. Diagnosis of HPB malignancy was made according to clinical, radiological and/or histological findings as dictated by center-specific protocols.

Categorical variables are expressed as numbers (*n*), with percentages in parenthesis, and continuous data as mean \pm standard deviation (SD) unless otherwise indicated. Statistical comparisons between groups were performed using Pearson’s Chi-squared test. Differences in the means and proportions between individual groups of continuous data were assessed using the independent samples t-test, following Levene’s test for equality of variances.¹⁹ A *p* value less than 0.05 was considered statistically significant.

Univariate and multivariable Cox proportional hazards models were fit to assess the impact of individual covariates on the instantaneous rate of clinical events, with time-to-event analysis ascertained through Kaplan-Meier estimates. Given that the development of IBD does not parallel that of PSC, the independent prognostic impact of IBD-phenotype was assessed separately as a time-fixed as well as a time-dependent covariate. All individual **covariates** were assessed for statistically significant interaction terms, **including patient demographic features**

(age and sex) and individual phenotypic descriptors for PSC and IBD subtypes separately. All analyses were stratified by geographical region (Australia, North America, Northern Europe, Central Europe, Western Europe or Southern Europe) and adjusted for year of PSC diagnosis. Incidence rates were calculated by the life tables' method. Statistical analyses were performed with IBM SPSS Statistics 22.0 (SPSS Inc, Chicago, IL).

Ethical approval

This study was conducted in accordance with the protocol and principles of the Declaration of Helsinki. The study protocol was reviewed and approved by the local institutional ethical boards of all participating centers.

RESULTS

Study population

We accrued clinical data pertaining to 7,931 patients (53,983 patient-years); however, those with inadequate follow-up or indeterminate diagnosis of PSC were exempted from further analysis (**Figure 1**). The final patient cohort consisted of **7,121** patients; either having PSC in its classical form (**89.8%**), as small-duct disease (**3.6%**), or the PSC/AIH-variant (6.6%) (**Table 1**). Observing the cohort in its entirety, the majority of patients were men (65.5%), with a mean age at diagnosis of 37 years versus 40 years in women ($p<0.001$). Seventy percent of all patients developed concomitant IBD prior to, at, or following PSC diagnosis; which under most circumstances was morphologically consistent with UC. However, **the development of UC** was less **common** in women than men (**48.1%** vs. **61.0%**, respectively; $p<0.001$), and in those with variant PSC sub-phenotypes relative to classical PSC (frequency of UC in patients with classical PSC: **58.1%** vs. **33.5%** in sdPSC, and vs. **47.7%** in PSC/AIH; $p<0.001$ for both pairwise comparisons) (**Supplementary Tables 1, 2 and 3**).

During the defined observation period, 20.2%, 37.0%, 52.3% and 63.6% of patients underwent liver transplantation or died at 5, 10, 15 and 20 years, respectively (**Figure 1**), yielding a median transplant-free survival time of **14.5** years (95% confidence interval [CI]: **13.6 – 15.2** years; **Figure 2A**). With regard to our secondary endpoint, 7.1%, 10.9%, 16.0% and 21.6% of the patient population developed a HPB malignancy at the aforementioned time points (**Figure 2B**) (overall $n = 721$).

The majority of HPB malignancy events were cholangiocarcinoma (CCA) ($n = 594$), and over one-third of all malignancies were detected in the first year following PSC diagnosis. The incidence of CCA increased with advancing age at PSC diagnosis (Supplementary Figure 1);

whilst hepatocellular carcinoma ($n = 59$) or gallbladder carcinoma ($n = 58$) were less frequent. Only ten patients across seven centers were diagnosed with pancreatic carcinoma. HPB malignancy developed most often in association with classical PSC, with only a small number of such events occurring in patients with sdPSC (1 CCA, 2 HCC, 1 pancreatic carcinoma) or PSC/AIH variants (12 CCA, 1 gallbladder carcinoma, 1 HCC). Overall, the development of HPB malignancy at any point during the clinical course was associated with a significantly increased risk of patient mortality (hazard ratio [HR]: 15.7, 95% CI: 14.12 – 17.34; $p < 0.001$).

Clinical stratifiers for liver transplantation/death and HPB malignancy

The incidence rates of clinical events according to baseline phenotypic descriptors are provided in **Supplementary Tables 4 and 5**. By univariate analysis, older age at diagnosis was associated with significantly poorer transplant-free survival; whereas female sex, CD (relative to UC), and sdPSC (relative to classical PSC) were identified as being protective (**Supplementary Table 6A**). No significant difference in transplant-free survival was observed between the PSC/AIH variant versus the classical PSC sub-phenotype (**Supplementary Figure 2A**), although patients with the former were at a low risk of developing HPB malignancy (**Supplementary Figure 2B**) (**Supplementary Table 6B**).

The number of patients with IBD increased during our observation period (from 3469 patients at baseline to 4985 patients by the end of our study). Given that intestinal disease onset did not necessarily parallel that in the liver, the impact of IBD was subsequently determined as a time-dependent covariate. In this context, both CD and an absence of IBD carried stratification properties of a lower risk PSC phenotype; whereas patients developing UC were at highest risk for disease progression, or future development of HPB malignancy (**Supplementary Table 6**).

Patient sex modifies the risk of liver disease progression in classical PSC

To verify the relative independence of predictive phenotypic features, a comparative multivariable evaluation was performed. Through multivariable Cox regression analysis the prognostic impact of advancing age at diagnosis, as well as protective influences of female sex, having **small duct disease**, or **CD at time of PSC diagnosis**, all retained statistical significance in terms of stratifying risk of liver disease progression (**Figures 3 and 4**).

Despite both factors being proven as independent risk-predictors, there was a statistically significant interaction ($p=0.013$) between patient sex and PSC sub-phenotypes when evaluating liver transplantation/death as an endpoint. To this effect, patients with sdPSC demonstrated significantly improved **transplant-free** survival, relative to same-sex counterparts with classical PSC and PSC/AIH, when matched for their age at PSC diagnosis as well as **baseline** IBD phenotype (**Figure 4A**). These differences were retained when adjusting for the latter as a time-dependent covariate in our multivariable analysis (**Table 2A**). Although women more commonly exhibited non-classical PSC sub-phenotypes than men, statistically significant differences in the risk of LTD between the sexes were retained when restricting our analyses to only those patients with classical PSC (**Table 2B**).

Unlike our primary endpoint, no statistically significant interactions were evident between patient sex and PSC sub-phenotypes when determining future HPB risk; wherein being female continued to exert a small, yet independent protective effect (but not an additive one) to that provided by small-duct disease (**Figures 3 and 4**) (**Table 3A and 3B**).

IBD phenotype as an independent predictor of clinical outcome in PSC

Crohn's disease (at time of PSC diagnosis) relative to UC continued to exert a protective influence with respect to transplant-free survival and the development of HPB malignancy,

irrespective of the effect exerted by sex and PSC sub-phenotype. Such impact was not demonstrated in the group without IBD at baseline (**Figure 4**). However, when addressing the impact of IBD as a time-dependent covariate, both CD and IBD-absence retained independent stratifying properties of a lower-risk PSC population (**Tables 2C** and **3C**). No statistically significant interactions existed between the different IBD phenotypes, and either PSC sub-phenotype or patient sex.

Reciprocally, development of UC prior to, or that which **manifest** during the clinical course of PSC, significantly increased the risk of LTD by **56%** and **15%** relative to CD or IBD-absence, respectively (**Table 2C**), and of HPB malignancy by approximately **45%** and **37%**, respectively (**Table 3C**). Of all patients with UC, **18.0 %** ($n = 718$) underwent colectomy before reaching a primary or secondary endpoint; however, no significant difference in outcome was evident in such individuals relative to those retaining an intact colon (HR for colectomy in terms of LTD and HPB malignancy: 0.90 (95% CI: 0.78 – 1.05; $p = 0.187$) and 0.81 (95% CI: 0.61 – 1.07; $p = 0.14$), respectively).

IBD phenotype overrides the prognostic impact of patient sex

The prognostic impact of IBD phenotype when assessed as a time-dependent variable negated the marginal protective influence of female sex. This means that although sex was an independent risk factor of both clinical endpoints statistically, there were no demonstrable differences **in either primary or secondary outcomes** between men and women when matched for IBD phenotype **as a time-dependent variable (data not shown)**. Moreover, the lower prevalence of UC in women (**Supplementary Table 1**) may account partially for differences in liver disease progression between the sexes.

DISCUSSION

PSC is a disease with significant clinical and societal burden, and in recognition of the hurdles involved in developing effective new therapies for patients, it is essential that robust descriptions of disease course are generated.^{2,3,4} In this study, we validate the critical importance of specific phenotypic variants (i.e., the more favourable prognosis that limited small-duct variants offers patients), the negative prognostic impact of ulcerative colitis on liver-related outcomes, and the high incidence of cholangiocarcinoma in the first year following PSC diagnosis.^{22,2} In addition, it is shown that patients with PSC and overlapping AIH-features carry a similar risk of liver disease progression to those with a more classical PSC phenotype; although development of HPB malignancy appears to be a rare event in PSC/AIH-overlap, and also for patients with a young presenting age at PSC diagnosis. Furthermore, we were able to address the prognostic impact of IBD development as a time-dependent covariate, recognising that development of UC is a key stratifier of adverse hepatobiliary consequences in PSC. Conversely, IBD-absence, and CD in particular, confer prognostic favour independent of the other phenotypic risk factors described.

To date, sex-specific variations in clinical phenotype and correlations with patient outcomes in PSC have lacked robust definition. Large scale studies have demonstrated the negative prognostic impact of male sex in patients with related disorders such as primary biliary cholangitis (PBC); specifically an association with treatment non-response and a higher incidence of HPB malignancy.^{23,24} As an immune-mediated disease PSC is somewhat atypical, with a propensity for ‘most’ patients being younger men. However, the sex-distribution of PSC appears more balanced if cholangiographic screening is applied to all IBD-patients irrespective of biochemical abnormalities or symptomatology.²⁵ In any event, utilising the large size of the

IPSCSG cohort, men with classical PSC are seen to carry a slight, albeit statistically significant increased risk of disease progression compared with women of matched phenotype.

Our analysis also demonstrates that women with PSC have a much lower prevalence of UC than men. **This is important because** IBD phenotype, particularly when determined as a time-dependent covariate, proves to be an independent risk factor for disease progression and **may explain the observed differences in outcome between sexes**. Conversely, patients without IBD or those having CD are at a comparatively lower risk of developing adverse events; a finding suggested previously in two single center studies, which we now validate convincingly.^{14,16} **Of note**, the IPSCSG has recently demonstrated genetic distinctions between patients with PSC and IBD versus those with IBD alone.^{26–28} Notwithstanding efforts to better understand clinical outcomes, **our** study further supports the need **to improve IBD classification in PSC**, particularly as the intestinal phenotype is often distinct compared to classical colitis descriptors,¹⁵ and more so given that genetic signals in PSC/CD may be disparate to those with PSC/UC.^{28,29} **Of note**, our study does not capture details pertaining to the precise distribution of intestinal inflammation; however, prior evidence suggests that CD in PSC is invariably localised to the colon, with isolated ileal disease being a seldom reported finding.^{14,16}

No significant outcome differences are apparent between men and women with the variant PSC sub-phenotypes, **and consequently** patients with sdPSC irrespective of gender experience a relatively sedentary clinical course compared with classical PSC. Perhaps more striking, however, is the highly similar transplant-free survival rate **seen for** patients with classical PSC **and** those with the PSC/AIH variant. Accepting the caveat that PSC/AIH lacks a codified diagnostic criteria,³⁰ these observations challenge the view of PSC/AIH variants imparting a lesser disease burden.³¹ Instead, our findings indicate that once overt sclerosing cholangitis has

manifest, liver disease may progress at a similar rate irrespective of the initial mode of disease presentation.

We also show how development of HPB malignancy (mainly CCA) manifests as a critical event in the clinical course of patients, particularly with advancing age at PSC diagnosis, and associated with significantly diminished patient survival. It is plausible that the reason for a third of CCA being identified within the first year following PSC diagnosis, is due to a delay in the latter's detection (length-time bias), and not being manifest until CCA is clinically overt. This observation highlights the need for improving CCA screening and surveillance, especially in high-risk PSC patients with coexisting UC. If better non-invasive surveillance methods for CCA surveillance became available, it could support the rationale for systematic screening for PSC in UC patients.²⁵ On the contrary, patients with small duct disease, perhaps indicative of PSC in an earlier form or of shorter duration, carry a lower risk of developing malignancy – as described previously.^{22,2} While this observation was somewhat expected, patients with the PSC/AIH-variant are also noted to develop HPB malignancy infrequently. This could possibly be a result of a lower UC burden,^{20,2,32,33} which as our data suggests, is itself an independent hazard for future carcinoma development. Furthermore, with only 10 cases during 51,500 patient years of follow-up we could not validate previous reports³⁷ of a significant increased incidence of pancreatic carcinomas, albeit accepting the clinical challenges that exist in differentiating distal cholangiocarcinomas from primary pancreatic lesions.

The natural history of PSC has previously been studied by some of the participating centers comprising the IPSCSG (Supplementary Table 7), although these cohorts are estimated to constitute, at most, <50% of our current patient population. Whilst certain patient characteristics that we describe mirror those in population-based registries,² ours is highly representative of a specialist-center PSC experience. In light of our prolonged study period,

transplant-center ‘designation’ and organ allocation policies have evolved significantly across institutions over time. Thus, it is not possible to accurately discriminate clinical outcomes based solely on the division between transplant versus non-transplant centers as conducted in other settings.² Admittedly, we do not present a population based epidemiological study, and due to the fact that more than 95% of included patients derived from centers with contemporary liver transplant activity, a degree of referral bias cannot be discounted. This may also explain the relatively low prevalence of sdPSC in our cohort.

Given the retrospective nature of our study, the interval frequency of repeated cholangiography varied between centers, therefore exhaustive surveillance imaging may not have been performed to exclude progression of all small duct cases to classical PSC. Similarly, there is no universally accepted guideline for repeated screening colonoscopy in those without IBD, hence we cannot discount that sub-clinical colitis may have developed in a subset of patients classified as having no IBD. Of note, our reported colectomy rate was 18% in patients with UC, which mirrors the incidence reported in single-center studies, but is lower than that observed in population-based cohorts and prospective multi-center registries of UC alone.^{34–36}

Our analyses were intentionally restricted to addressing the prognostic impact of well-defined patient phenotypes. Consequently data pertaining to laboratory variables, extent of strictures, intervals of surveillance imaging or specific pharmacological interventions (e.g. ursodeoxycholic acid and/or immunosuppression) fell outside of the current study’s remit. Further large-scale investigation of therapeutic impact is of critical importance, given the inconsistently reported effects of these agents on disease progression and malignancy risk in PSC.⁸ Additionally, as a systematic autopsy review was not performed from all mortality cases it is plausible that the incidence of HPB malignancy may in fact be higher than actually reported,³⁷ particularly as CCA cannot always be discriminated from more benign changes in

PSC.³⁸ We are also unable to classify all causes of death in our retrospective patient cohort, although previous studies indicate that mortality in PSC is invariably due to liver disease or a complication of coexisting IBD.^{2,39} A further restriction due to the retrospective nature and prolonged follow-up period (since 1980) is the fact that serum IgG4-levels were not determined systematically in all patients. Therefore it is not possible to conclusively exclude IgG4 associated cholangiopathy within a subset of our population.

The IPSCSG study confirms significant **phenotypic** diversity across the global PSC patient population. **The estimates provided for transplant-free survival and the lifetime risk of HPB malignancy, would facilitate** appropriate patient counselling and also aid in the future evaluation of potential new approaches to malignancy screening. In a drive to limit heterogeneity in clinical trials, which currently group together individuals at a high-risk of disease progression (classical PSC and UC) **together with patients at intermediate risk (CD or IBD-absence) and low risk (sdPSC),** our data underpins a collaborative effort to better appraise future therapeutic ventures for this orphan disease. As a clear consequence of our findings, future clinical trials may now be able to stratify entry according to a combination of precise phenotypic risk factors, limit the heterogeneity within studied cohorts, and provide a more objective evaluation of therapeutic efficacy in specific patient groups.

Table 1: Summary of Patient Characteristics

No. of pts.	7121
No. of men	4661 (65.5%)
Age at diagnosis:	
- Mean	38.5 yrs. (SD: 15.5)
- ≤ 20 yrs.	940 (13.2%)
- 21 – 30 yrs.	1508 (21.2%)
- 31 – 40 yrs.	1617 (22.7%)
- 41 – 50 yrs.	1435 (20.2%)
- 51 – 60 yrs.	953 (13.4%)
- > 60 yrs.	665 (9.3%)
- <i>unknown</i>	3 (0.04%)
PSC sub-phenotype:	
- classical PSC	6397 (89.8%)
- small duct PSC	254 (3.6%)
- PSC / AIH variant	470 (6.6%)
Diagnosis year:	
- 1980 – 1984	217 (3.0%)
- 1985 – 1989	424 (6.0%)
- 1990 – 1994	773 (10.9%)
- 1995 – 1999	1414 (19.9%)
- 2000 – 2004	1802 (25.3%)
- 2005 – 2010	2491 (35.0%)
IBD phenotype at baseline:	
- ulcerative colitis	2761 (38.8%)
- Crohn's disease	595 (8.4%)
- indeterminate colitis	113 (1.6%)
- no IBD	3082 (43.3%)
- unknown timing	503 (7.1%)
- unknown IBD status	67 (0.9%)
IBD phenotype at end of follow-up:	
- ulcerative colitis	3989 (56.0%)
- Crohn's disease	786 (11.0%)
- indeterminate colitis	210 (2.9%)
- no IBD	2069 (29.1%)
- unknown IBD status	67 (0.9%)

Table 2: Risk Stratification of Liver Transplantation / Death by Disease Phenotype

		Reference phenotype	Adjusted hazard ratio (95% CI)	p-value
A) PSC phenotype	<i>Male</i>			
	Small-duct PSC	vs Classical PSC	0.23 (0.13 – 0.40)	<0.001
	PSC/AIH variant	vs Classical PSC	0.73 (0.56 – 0.94)	0.015
	PSC/AIH variant	vs Small-duct PSC	3.18 (1.71 – 5.92)	<0.001
	<i>Female</i>			
	Small-duct PSC	vs Classical PSC	0.48 (0.29 – 0.77)	0.003
B) Sex	<i>Classical PSC</i>			
	Female	vs Male	0.84 (0.77 – 0.92)	0.022
	<i>Small-duct PSC</i>			
	Female	vs Male	1.76 (0.84 – 3.69)	0.13
	<i>PSC/AIH variant</i>			
	Female	vs Male	1.38 (0.97 – 1.97)	0.075
C) IBD phenotype	Crohn's disease	vs Ulcerative colitis	0.64 (0.54 – 0.75)	<0.001
	Indeterminate colitis	vs Ulcerative colitis	0.94 (0.71 – 1.26)	0.69
	No IBD	vs Ulcerative colitis	0.87 (0.79 – 0.95)	0.002
	Crohn's disease	vs no IBD	0.73 (0.62 – 0.87)	<0.001
	Indeterminate colitis	vs no IBD	1.10 (0.83 – 1.48)	0.51
	Indeterminate colitis	vs Crohn's disease	1.50 (1.09 – 2.07)	0.013

* All analyses are stratified by geographical region of diagnosis; adjusted for calendar year and age at diagnosis. Inflammatory bowel disease phenotype is defined as a time dependent covariate. Hazard ratios for PSC sub-phenotypes are presented separately for men and women, and separately for sex are presented separately for PSC sub-phenotype, given the presence of a significant interaction term between gender and PSC sub-phenotype ($p = 0.005$).

Table 3: Stratification of Hepatopancreatobiliary Malignancy Risk by Disease Phenotype

		Reference phenotype	Adjusted hazard ratio (95% CI)	p-value
A) PSC phenotype	Small-duct PSC	vs Classical PSC	0.19 (0.07 – 0.51)	0.001
	PSC/AIH variant	vs Classical PSC	0.31 (0.17 – 0.55)	<0.001
	PSC/AIH variant	vs Small-duct PSC	1.62 (0.52 – 5.04)	0.41
B) Sex	Female	vs Male	0.68 (0.57 – 0.82)	0.001
C) IBD phenotype	Crohn's disease	vs Ulcerative colitis	0.69 (0.52 – 0.92)	0.01
	Indeterminate colitis	vs Ulcerative colitis	1.03 (0.52 – 1.75)	0.931
	No IBD	vs Ulcerative colitis	0.73 (0.61 – 0.87)	<0.001
	Crohn's disease	vs no IBD	0.96 (0.71 – 1.29)	0.77
	Indeterminate colitis	vs no IBD	1.41 (0.82 – 2.44)	0.22
	Indeterminate colitis	vs Crohn's disease	1.48 (0.82 – 2.67)	0.20

* All analyses stratified by geographical region of diagnosis; adjusted for calendar year and age at diagnosis. Inflammatory bowel disease phenotype is defined as a time dependent covariate.

Figure 1: Study cohort

At time of analysis data were available for 7,931 patients. However, following exclusion of groups with an alternate diagnose or inadequate follow-up, the final study group consisted of 7,121 patients of which 2,616 underwent liver transplantation or died, with a total of 721 developing primary hepatopancreatobiliary malignancy.

Figure 2: Cumulative incidence of clinical events

Kaplan-Meier estimates of [A] liver transplant (LT)-free survival rate across the patient population; and [B] incidence of all hepatopancreatobiliary (HPB) malignancies. Notably, 37.8% ($n = 272$) of all HPB malignancies occurred in the first year of PSC diagnosis, with the vast majority being cholangiocarcinoma during this time (incidence rate in the first year after PSC diagnosis: 2.6 cases per-100 patient-years).

Patients with unknown transplantation, mortality or malignancy status at time of study completion were excluded from respective analysis.

Figure 3: Impact of Patient Age and Gender on Clinical Outcome

Cox plots with regard to liver transplantation (LT) or hepatopancreatobiliary (HPB) malignancy. All data are stratified by geographical region of referring center and year of diagnosis, presented according to patient age at diagnosis and weighted for patient gender, inflammatory bowel disease (IBD) phenotype at baseline, and PSC sub-phenotype [A + B]; or patient gender weighted for patient age at diagnosis, IBD phenotype at baseline, and PSC sub-phenotype [C + D].

Figure 4: Impact of Variant PSC Sub-phenotypes and IBD Phenotypes on Clinical Outcome

Cox plots with regard to liver transplantation (LT) or hepatopancreatobiliary (HPB) malignancy. All data are stratified by geographical region of referring center and year of diagnosis, presented according to PSC sub-phenotype weighted for patient age at PSC diagnosis, gender, and inflammatory bowel disease (IBD) phenotype at baseline [A + B]; or patient IBD phenotype at baseline weighted for age at PSC diagnosis, gender, and PSC sub-phenotype [C + D].

Contributors:

TJW, PJT, BEH and KMB contributed equally to the manuscript and were primarily involved in data collection, validation, analysis, and manuscript preparation. AB, GMH, THK and CPS contributed to the data analysis and the manuscript preparation. BEH was the official study statistician who conducted and supervised statistical analysis and data interpretation. TJW, PJT, MI, HL, CYP, KH, DG, MAF, H-UM, DT, RKW, JF, TM, OC, KS, KNL, SA, SPP, CL, AM, SN, CLB, AF, EH, KKY, PM, UB, DKH, AP, CNM, GND, BE, PI, CPB, GIK, CS, VZ, LF, FB, MM, BDJ, KS, CR, KJ, MBdV, FS, **AC**, MT, THK, ES, MM, CPS, KDL, GMH and KMB were all involved in patient recruitment and assembling individual center data. All authors read and approved the final manuscript before submission.

Declaration of interests

All authors declare no competing interests regarding this study.

Acknowledgements

TJW was supported by the German Federal Ministry of Education and Research through the Integrated Research and Treatment Center Transplantation (reference number 01EO0802) at Hannover Medical School. PJT has received funding from the Wellcome Trust. PJT and GMH also receive funding and salary support from the NIHR Birmingham Biomedical Research Unit (- the views expressed are those of the authors and not necessarily those of the NHS, the NIHR or the Department of Health, UK). SPP was supported by the NIHR University College London Hospitals BRC. KNL was supported by NIH R01 DK84960. TM was supported by the German Research Community, Grants MU 2864/1-1 and MU 2864/1-3. CS was supported by the German Research Community (DFG, SFB 841), by the YAEL-Foundation and the Helmut and Hannelore Greve Foundation

The following collaborators are acknowledged for their contributions with data acquisition:

Benoît Almer, MD (University of Lund)

The IPSCSG would also like to extend thanks to all patients who contributed to data in the study, as well as the patient support groups (PSC partners and PSC support).

REFERENCES

- 1 Hirschfield GM, Karlsen TH, Lindor KD, Adams DH. Primary sclerosing cholangitis. *Lancet* 2013; **382**: 1587–99.
- 2 Boonstra K, Weersma RK, van Erpecum KJ, *et al.* Population-based epidemiology, malignancy risk, and outcome of primary sclerosing cholangitis. *Hepatology* 2013; **58**: 2045–55.
- 3 Boonstra K, Beuers U, Ponsioen CY. Epidemiology of primary sclerosing cholangitis and primary biliary cirrhosis: A systematic review. *J Hepatol* 2012; **56**: 1181–8.
- 4 Lindkvist B, Benito de Valle M, Gullberg B, Björnsson E. Incidence and prevalence of primary sclerosing cholangitis in a defined adult population in Sweden. *Hepatology* 2010; **52**: 571–7.
- 5 Patkowski W, Skalski M, Zieniewicz K, Nyckowski P, Smoter P, Krawczyk M. Orthotopic liver transplantation for cholestatic diseases. *Hepatogastroenterology* 2010; **57**: 605–10.
- 6 Charman S, Copley L, Tovikkai C, Meulen J van der. UK Liver Transplant Audit (NHS Blood and Transplant). 2012. <https://www.rcseng.ac.uk/surgeons/research/surgical-research/docs/liver-transplant-audit-report-2012>.
- 7 Fosby B, Melum E, Bjørø K, *et al.* Liver transplantation in the Nordic countries - An intention to treat and post-transplant analysis from The Nordic Liver Transplant Registry 1982-2013. *Scand J Gastroenterol* 2015; **50**: 797–808.
- 8 Lindor KD, Kowdley KV, Luketic VAC, *et al.* High-dose ursodeoxycholic acid for the treatment of primary sclerosing cholangitis. *Hepatology* 2009; **50**: 808–14.

- 9 Trivedi PJ, Corpechot C, Pares A, Hirschfield GM. Risk stratification in autoimmune cholestatic liver diseases: Opportunities for clinicians and trialists. *Hepatology* 2016; **63**: 644–59.
- 10 EASL Clinical Practice Guidelines: management of cholestatic liver diseases. *J Hepatol* 2009; **51**: 237–67.
- 11 Chapman R, Fevery J, Kalloo A, *et al.* Diagnosis and management of primary sclerosing cholangitis. *Hepatology* 2010; **51**: 660–78.
- 12 Lindor KD, Kowdley KV, Harrison ME. ACG Clinical Guideline: Primary Sclerosing Cholangitis. *Am J Gastroenterol* 2015; **110**: 646–59.
- 13 Boberg KM, Chapman RW, Hirschfield GM, Lohse AW, Manns MP, Schrumpf E. Overlap syndromes: The International Autoimmune Hepatitis Group (IAIHG) position statement on a controversial issue. *J Hepatol* 2011; **54**: 374–85.
- 14 Halliday JS, Djordjevic J, Lust M, *et al.* A unique clinical phenotype of primary sclerosing cholangitis associated with Crohn’s disease. *J Crohns Colitis* 2012; **6**: 174–81.
- 15 Boonstra K, van Erpecum KJ, van Nieuwkerk KMJ, *et al.* Primary sclerosing cholangitis is associated with a distinct phenotype of inflammatory bowel disease. *Inflamm Bowel Dis* 2012; **18**: 2270–6.
- 16 Fevery J, Van Steenberghe W, Van Pelt J, *et al.* Patients with large-duct primary sclerosing cholangitis and Crohn’s disease have a better outcome than those with ulcerative colitis, or without IBD. *Aliment Pharmacol Ther* 2016; **43**: 612–20.
- 17 Dignass A, Eliakim R, Magro F, *et al.* Second European evidence-based consensus on the diagnosis and management of ulcerative colitis part 1: definitions and diagnosis. *J Crohns Colitis* 2012; **6**: 965–90.

- 18 Van Assche G, Dignass A, Panes J, *et al.* The second European evidence-based Consensus on the diagnosis and management of Crohn's disease: Definitions and diagnosis. *J Crohns Colitis* 2010; **4**: 7–27.
- 19 Levene H. Contributions to Probability and Statistics: Essays in Honor of Harold Hotelling. Stanford University Press, 1960.
- 20 Boberg KM, Bergquist A, Mitchell S, *et al.* Cholangiocarcinoma in primary sclerosing cholangitis: risk factors and clinical presentation. *Scand J Gastroenterol* 2002; **37**: 1205–11.
- 21 Rudolph G, Gotthardt DN, Kloeters-Plachky P, Kulaksiz H, Schirmacher P, Stiehl A. In PSC with colitis treated with UDCA, most colonic carcinomas develop in the first years after the start of treatment. *Dig Dis Sci* 2011; **56**: 3624–30.
- 22 Björnsson E, Olsson R, Bergquist A, *et al.* The Natural History of Small-Duct Primary Sclerosing Cholangitis. *Gastroenterology* 2008; **134**: 975–80.
- 23 **Carbone M, Mells GF, Pells G, et al.** Sex and age are determinants of the clinical phenotype of primary biliary cirrhosis and response to ursodeoxycholic Acid. *Gastroenterology* 2013; **144**: 560–9.e7.
- 24 **Trivedi PJ, Lammers WJ, van Buuren HR, et al.** Stratification of hepatocellular carcinoma risk in primary biliary cirrhosis: a multicentre international study. *Gut* 2016; **65**: 321–9.
- 25 Lunder AK, Hov JR, Borthne A, *et al.* Prevalence of Sclerosing Cholangitis Detected by Magnetic Resonance Cholangiography in Patients With Long-term Inflammatory Bowel Disease. *Gastroenterology* 2016; **151**: 660–9.e4.
- 26 **Ellinghaus D, Folseraas T, Holm K, et al.** Genome-wide association analysis in primary sclerosing cholangitis and ulcerative colitis identifies risk loci at GPR35 and TCF4. *Hepatology* 2013; **58**: 1074–83.

- 27 **Liu JZ, Hov JR, Folseraas T, et al.** Dense genotyping of immune-related disease regions identifies nine new risk loci for primary sclerosing cholangitis. *Nat Genet* 2013; **45**: 670–5.
- 28 **Ji S-G, Juran BD, Mucha S, et al.** Genome-wide association study of primary sclerosing cholangitis identifies new risk loci and quantifies the genetic relationship with inflammatory bowel disease. *Nat Genet* 2016; **advance online publication**. DOI:10.1038/ng.3745.
- 29 Maroni L, van de Graaf SFJ, Hohenester SD, Oude Elferink RPJ, Beuers U. Fucosyltransferase 2: A Genetic Risk Factor for Primary Sclerosing Cholangitis and Crohn's Disease—A Comprehensive Review. *Clin Rev Allergy Immunol* 2015; **48**: 182–91.
- 30 Trivedi PJ, Hirschfield GM. Review article: overlap syndromes and autoimmune liver disease. *Aliment Pharmacol Ther* 2012; **36**: 517–33.
- 31 Floreani A, Rizzotto ER, Ferrara F, et al. Clinical Course and Outcome of Autoimmune Hepatitis/Primary Sclerosing Cholangitis Overlap Syndrome. *Am J Gastroenterol* 2005; **100**: 1516–22.
- 32 Rudolph G, Gotthardt D, Kloeters-Plachky P, Rost D, Kulaksiz H, Stiehl A. In PSC with dominant bile duct stenosis, IBD is associated with an increase of carcinomas and reduced survival. *J Hepatol* 2010; **53**: 313–7.
- 33 Gulamhusein AF, Eaton JE, Tabibian JH, Atkinson EJ, Juran BD, Lazaridis KN. Duration of Inflammatory Bowel Disease Is Associated With Increased Risk of Cholangiocarcinoma in Patients With Primary Sclerosing Cholangitis and IBD. *Am J Gastroenterol* 2016; published online March 22. DOI:10.1038/ajg.2016.55.

- 34 Manetti N, Bagnoli S, Rogai F, *et al.* Disease Course and Colectomy Rate of Ulcerative Colitis: A Follow-up Cohort Study of a Referral Center in Tuscany. *Inflamm Bowel Dis* 2016; **22**: 1945–53.
- 35 Samuel S, Ingle SB, Dhillon S, *et al.* Cumulative incidence and risk factors for hospitalization and surgery in a population-based cohort of ulcerative colitis. *Inflamm Bowel Dis* 2013; **19**: 1858–66.
- 36 Rutter MD, Saunders BP, Wilkinson KH, *et al.* Thirty-year analysis of a colonoscopic surveillance program for neoplasia in ulcerative colitis. *Gastroenterology* 2006; **130**: 1030–8.
- 37 Bergquist A, Ekbom A, Olsson R, *et al.* Hepatic and extrahepatic malignancies in primary sclerosing cholangitis. *J Hepatol* 2002; **36**: 321–7.
- 38 Charatcharoenwitthaya P, Enders FB, Halling KC, Lindor KD. Utility of serum tumor markers, imaging, and biliary cytology for detecting cholangiocarcinoma in primary sclerosing cholangitis. *Hepatology* 2008; **48**: 1106–17.
- 39 Claessen MMH, Vleggaar FP, Tytgat KMAJ, Siersema PD, van Buuren HR. High lifetime risk of cancer in primary sclerosing cholangitis. *J Hepatol* 2009; **50**: 158–64.

Figure 1





Figure 2

A



B




Figure 3





Figure 4



Supplementary Table 1: Patient Characteristics by Gender *

A) Demographics and phenotype **		
	Men (n = 4661)	Women (n = 2454) ***
Age at diagnosis: ****		
- Mean	37 yrs. (SD: 15)	40 yrs. (SD: 16)
- <= 20 yrs.	660 (14.2%)	278 (11.4%)
- 21 – 30 yrs.	1065 (22.8%)	442 (18.0%)
- 31 – 40 yrs.	1084 (23.3%)	532 (21.7%)
- 41 – 50 yrs.	904 (19.4%)	531 (21.7%)
- 51 – 60 yrs.	550 (11.8%)	403 (16.4%)
- > 60 yrs.	397 (8.5%)	266 (10.8%)
PSC sub-phenotype: ****		
- classical PSC	4231 (90.8%)	2160 (88.0%)
- small duct PSC	158 (3.4%)	96 (3.9%)
- PSC / AIH variant	271 (5.8%)	198 (8.1%)
Diagnosis year:		
- 1980 – 1984	144 (3.1%)	73 (3.0%)
- 1985 – 1989	304 (6.5%)	120 (4.9%)
- 1990 – 1994	524 (11.2%)	248 (10.1%)
- 1995 – 1999	937 (20.1%)	477 (19.4%)
- 2000 – 2004	1176 (25.2%)	623 (25.4%)
- 2005 – 2010	1576 (33.8%)	913 (37.2%)
IBD phenotype at baseline: ****		
- ulcerative colitis	1935 (45.4%)	823 (36.0)
- Crohn’s disease	362 (8.5%)	233 (9.5)
- indeterminate colitis	76 (1.8%)	37 (1.6)
- no IBD	1890 (44.3%)	1190 (52.1)
IBD phenotype at end of follow-up: ****		
- ulcerative colitis	2818 (61.0)	1168 (48.1)
- Crohn’s disease	466 (10.1)	318 (13.1)
- indeterminate colitis	143 (3.1)	67 (2.8)
- no IBD	1193 (25.5)	874 (36.0.7)
B) Clinical events ****	Incidence rate per-100-pt. yrs. (95%. C.I.)	
Liver transplantation or death	5.58 (5.34-5.82)	5.16 (4.83-5.48)
Hepatopancreatobiliary malignancy		
- overall	1.55 (1.41-1.68)	1.10 (0.94-1.25)
- cholangiocarcinoma	1.28 (0.86-1.71)	0.90 (0.43-1.37)

* Data presented as absolute number (%) unless otherwise indicated.
** Data presented only for patients in whom complete respective data are available.
*** Six patients did not have gender data documented
**** Indicates statistically significant differences of covariate frequency between all subgroups listed ($p < 0.05$).

Supplementary Table 2: Patient Characteristics by PSC Sub-phenotype *

A) Demographics and phenotype **			
	Classical PSC (n= 6397)	Small-duct PSC (n = 254)	PSC / AIH variant (n = 470)
No. of men	4232 (66.2%)	158 (62.2%)	271 (57.8%)
Age at diagnosis:			
- Mean	39 yrs. (SD: 15.4)	37yrs. (SD: 14.8)	32 yrs. (SD: 15)
- < 20 yrs.	779 (12.2%)	35 (13.8%)	126 (26.8%)
- 21 – 30 yrs.	1323 (20.7%)	59 (23.2%)	126 (26.8%)
- 31 – 40 yrs.	1456 (22.8%)	68 (26.8%)	93 (19.8%)
- 41 – 50 yrs.	1327 (20.8%)	43 (16.9%)	65 (13.8%)
- 51 – 60 yrs.	884 (13.8%)	32 (12.6%)	37 (7.9%)
- > 60 yrs.	625 (9.8%)	17 (6.7%)	23 (4.9%)
Diagnosis year:			
- 1980 – 1984	213 (3.3%)	2 (0.8%)	2 (0.4%)
- 1985 – 1989	404 (6.3%)	9 (3.5%)	11 (2.3%)
- 1990 – 1994	723 (11.3%)	18 (7.1%)	32 (6.8%)
- 1995 – 1999	1287 (20.1%)	47 (18.5%)	80 (17.0%)
- 2000 – 2004	1603 (25.1%)	79 (31.1%)	120 (25.5%)
- 2005 – 2010	2167 (33.9%)	99 (39.0%)	225 (47.9%)
IBD phenotype at baseline:			
- ulcerative colitis	2535 (43.2%)	67 (27.9%)	159 (36.2%)
- Crohn's disease	545 (9.3%)	24 (10.0%)	26 (5.9%)
- indeterminate colitis	98 (1.7%)	6 (2.5%)	9 (2.1%)
- no IBD	2694 (45.9%)	143 (59.6%)	245 (55.8%)
IBD phenotype at end of study:			
- ulcerative colitis	3682 (58.1%)	85 (33.5%)	222 (47.7%)
- Crohn's disease	718 (11.3%)	30 (11.8%)	38 (8.2%)
- indeterminate colitis	185 (2.9%)	7 (2.8%)	18 (3.9%)
- no IBD	1750 (27.6%)	132 (52.0%)	187 (40.2%)
B) Clinical events **	Incidence rate per-100-pt. yrs. (95%. C.I.)		
Liver transplantation or death	5.62 (5.42 -5.83)	2.32 (1.67 – 3.00)	4.70 (3.97 – 5.43)
Hepatopancreatobiliary malignancy			
-overall	1.52 (1.41 –1.63)	0.20 (0.00 –0.39)	0.43 (0.20 – 0.65)
- cholangiocarcinoma	1.25 (0.90–1.60)	No cases	0.37 (0.16 – 0.58)

* Data presented as absolute number (%) unless otherwise indicated.

** Data presented only for patients in whom complete respective data are available.

Supplementary Table 3: Patient Characteristics by IBD phenotype (at baseline) *

A) Demographics and phenotype **				
	Ulcerative colitis (n = 2761)	Crohn's Disease (n= 595)	Indeterminate (n = 113)	No IBD (n = 3082)
No. of men	1935 (70.2)	362 (60.8)	76 (67.3)	1890 (61.4)
Age at diagnosis:				
- Mean	37 yrs. (SD: 15)	38 yrs. (SD: 16)	35 yrs. (SD: 14)	40 yrs. (SD: 16)
- ≤ 20 yrs.	410 (14.8%)	91 (15.3%)	17 (15.0%)	350 (11.4%)
- 21 – 30 yrs.	646 (23.4%)	125 (21.0%)	36 (31.9%)	585 (19.0%)
- 31 – 40 yrs.	671 (24.3%)	136 (22.9%)	24 (21.2%)	660 (21.4%)
- 41 – 50 yrs.	510 (18.5%)	116 (19.5%)	17 (15.0%)	664 (21.6%)
- 51 – 60 yrs.	336 (12.2%)	74 (12.4%)	13 (11.5%)	452 (14.7%)
- > 60 yrs.	188 (6.8%)	53 (8.9%)	6 (5.3%)	368 (12.0%)
PSC sub-phenotype:				
- classical PSC	2535 (91.8%)	545 (91.6%)	98 (86.7%)	2694 (87.4%)
- small duct PSC	67 (2.4%)	24 (4.0%)	6 (5.3%)	143 (4.6%)
- PSC / AIH variant	159 (5.8%)	26 (4.4%)	9 (8.0%)	245 (7.9%)
Diagnosis year:				
- 1980 – 1984	75 (2.7%)	9 (1.5%)	4 (3.5%)	91 (3.0%)
- 1985 – 1989	166 (6.0%)	23 (3.9%)	6 (5.3%)	167 (5.4%)
- 1990 – 1994	327 (11.8%)	41 (6.9%)	16 (14.2%)	299 (9.7%)
- 1995 – 1999	561 (20.3%)	104 (17.5%)	15 (13.3%)	620 (20.1%)
- 2000 – 2004	705 (25.5%)	165 (27.7%)	27 (23.9%)	783 (25.4%)
- 2005 – 2010	927 (33.6%)	253 (42.5%)	45 (39.8%)	1122 (36.4%)
B) Clinical events **	Incidence rate per-100-pt. yrs. (95% C.I.)			
Liver transplantation or death	5.36 (5.06-5.67)	3.89 (3.30-4.47)	4.47 (3.07-5.88)	5.82 (5.51-6.13)
Hepatopancreatobiliary malignancy				
-overall	1.48 (1.31-1.64)	1.21 (0.88-1.55)	1.43 (0.62-2.24)	1.34 (1.19-1.50)
- cholangiocarcinoma	1.22 (0.72-1.72)	1.02 (0.03-2.02)	1.19 (0.00-3.07)	1.11 (0.60-1.62)

* Data presented as absolute number (%) unless otherwise indicated.

** Data presented only for patients in whom complete respective data are available.

Supplementary Table 4: Incidence Rates (IR) per-100-pt. yrs. of Liver transplantation / Death According to Phenotype

Event: liver transplantation / death								
	Male				Female			
	<i>UC</i>	<i>CD</i>	<i>IC</i>	<i>No-IBD</i>	<i>UC</i>	<i>CD</i>	<i>IC</i>	<i>No-IBD</i>
<i>Classical PSC</i>								
IR:	5.5	4.3	4.6	6.3	5.3	3.4	5.5	5.7
1y survival:	94%	96%	97%	92%	95%	96%	100%	94%
5y survival:	77%	80%	82%	71%	79%	85%	73%	77%
10y survival:	59%	67%	73%	55%	61%	72%	62%	60%
20y survival:	30%	52%	37%	31%	23%	67%	40%	35%
<i>sdPSC</i>								
IR:	2.5	0.0	0.0	2.2	2.7	4.0	0.0	2.5
1y survival:	96%	100%	100%	99%	100%	100%	100%	95%
5y survival:	96%	100%	100%	89%	100%	88%	100%	86%
10y survival:	96%	100%	100%	89%	75%	88%	-	80%
20y survival:	84%	-	-	82%	56%	-	-	67%
<i>PSC/AIH-overlap</i>								
IR:	4.1	4.8	2.1	3.9	5.2	6.6	0.0	5.5
1y survival:	96%	100%	100%	96%	97%	92%	100%	96%
5y survival:	86%	92%	83%	78%	79%	61%	-	81%
10y survival:	73%	69%	83%	68%	69%	41%	-	56%
20y survival:	45%	69%	-	55%	30%	41%	-	29%

Supplementary Table 5: Incidence Rates (IR) per-100-pt. yrs. of HPB malignancy
According to Phenotype

Event: hepatopancreatobiliary (HPB) malignancy *								
	Male				Female			
	<i>UC</i>	<i>CD</i>	<i>IC</i>	<i>No-IBD</i>	<i>UC</i>	<i>CD</i>	<i>IC</i>	<i>No-IBD</i>
<i>Classical PSC</i>								
IR; 1 st yr. only:	3.1	2.2	3.5	3.8	2.2	2.1	1.9	2.6
IR; overall:	1.6	1.6	1.4	1.7	1.5	0.6	1.5	1.1
1y survival:	96%	97%	95%	94%	97%	97%	97%	96%
5y survival:	92%	92%	93%	90%	92%	96%	91%	92%
10y survival:	86%	87%	93%	86%	86%	95%	78%	90%
20y survival:	70%	73%	82%	75%	68%	95%	78%	83%
<i>sdPSC</i>								
IR; 1 st yr. only:	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
IR; overall:	0.0	0.0	0.0	0.0	0.4	0.0	0.0	0.5
1y survival:	100%	100%	100%	100%	100%	100%	100%	100%
5y survival:	100%	100%	100%	100%	100%	100%	100%	98%
10y survival:	100%	100%	100%	100%	89%	100%	-	92%
20y survival:	100%	-	-	100%	89%	-	-	92%
<i>PSC/AIH-overlap</i>								
IR; 1 st yr. only:	1.5	6.5	0.0	0.7	0.0	0.0	0.0	0.8
IR; overall:	0.7	2.0	0.0	0.2	0.2	1.2	0.0	0.1
1y survival:	96%	92%	100%	99%	100%	100%	100%	99%
5y survival:	94%	81%	100%	98%	98%	89%	-	99%
10y survival:	94%	81%	100%	98%	98%	89%	-	99%
20y survival:	94%	81%	-	98%	98%	-	-	99%

* For HPB malignancy, IR are provided for events in the 1st year only as well as overall

Supplementary Table 6: Univariate Risk Factors for Disease Progression *

Risk factor	Crude Hazard Ratio (95% C.I.)	p value
A) Liver transplantation / death		
Age at diagnosis **	1·022 (1·019 – 1·025)	< 0·0001
Gender		
Male	1 (reference)	
Female	0·88 (0·81 – 0·96)	0·002
PSC sub-phenotype		
- classical PSC	1 (reference)	
- small duct PSC	0·30 (0·21 – 0·42)	< 0·001
- PSC / AIH variant	0·81 (0·68 – 0·96)	0·015
IBD phenotype (<i>baseline</i>)		
- ulcerative colitis	1 (reference)	
- Crohn's disease	0·64 (0·53 – 0·76)	<0·0001
- indeterminate	0·86 (0·61 – 1·22)	0·40
- no IBD	1·01 (0·93 – 1·10)	0·89
IBD phenotype (<i>prior-to-endpoint</i>) ***		
- ulcerative colitis	1 (reference)	
- Crohn's disease	0·62 (0·52 – 0·72)	< 0·001
- indeterminate	0·91 (0·68 – 1·21)	0·52
- no IBD	0·90 (0·83 – 0·99)	0·03
B) Hepatopancreatobiliary malignancy		
Age at diagnosis **	1·03 (1·03 – 1·04)	< 0·001
Gender		
Male	1 (reference)	
Female	0·68 (0·57 – 0·80)	< 0·001
PSC biliary phenotype		
- classical PSC	1 (reference)	
- small duct PSC	0·15 (0·06 – 0·40)	< 0·001
- PSC / AIH variant	0·26 (0·15 – 0·44)	< 0·001
IBD phenotype (<i>baseline</i>)		
- ulcerative colitis	1 (reference)	
- Crohn's disease	0·73 (0·54 – 0·96)	0·04
- indeterminate	1·09 (0·61 – 1·94)	0·77
- no IBD	0·88 (0·75 – 1·04)	0·14
IBD phenotype (<i>prior-to-endpoint</i>) ***		
ulcerative colitis	1 (reference)	
- Crohn's disease	0·68 (0·51 – 0·91)	0·008
- indeterminate	0·94 (0·55 – 1·61)	0·82
- no IBD	0·77 (0·65 – 0·92)	0·004

*All analysis stratified by geographical region of participating centre and adjusted by patient year of diagnosis.

** Per 1-yr. increase in age.

*** Assessed as a time-dependent covariate

Supplementary Table 7: Previously published clinical outcome studies in PSC *

Geographical location	Study type	Study period or last reported follow-up date – previously reported	Maximum No. pts. – previously reported
<i>Multi-national</i>			
Italy, Norway, Spain, Sweden, UK	Observational	1998 ^{1,2}	394
Scandinavia	Clinical trial	2009 ^{3–5}	219 **
Finland, the Netherlands, Norway, UK	Investigative biomarker	2012 ⁶	305
Germany and Sweden	Observational	1989 – 2008 (Germany) ^{7,8} 1992 – 2005 (Sweden)	345
Germany and Norway	Observational	2014 ¹¹	638
Germany and Norway	Investigative biomarker	2006 – 2015 (Germany) ¹² 2008 – 2012 (Norway)	318
<i>Belgium</i>			
Leuven	Observational	1975 – 2012 ^{13,14}	240
<i>Canada</i>			
Toronto, ON	Observational	2009 ¹⁵	168
<i>France</i>			
Paris	Observational	2008 ¹⁶	150
<i>Germany</i>			
Heidelberg	Observational / investigative biomarker	2012 ^{17–21}	281 ***
Hannover	Observational	2006 ¹⁰	273
Hamburg and Hannover	Observational	2013 ⁹	509
<i>Italy</i>			
Multi-regional	Observational	1994 ²²	117
<i>The Netherlands</i>			
Multi-regional	Observational	2008 ^{23–27}	590 ***
<i>Sweden</i>			
Multi-regional	Observational	1992 ²⁸	305
Stockholm	Observational	1970 – 2004 ^{29–31}	604
<i>USA</i>			
Multi-regional	Clinical trial	2009 ^{32–34}	150
Multi-regional	Observational	1995 – 2005 ³³	784
Minnesota	Observational	1970 – 1997 ^{36,37}	174
California	Observational	2000 – 2006 ³⁸	169
<i>UK</i>			
London	Observational	2011 ³⁹	128
London	Observational	1990 – 2009 ⁴⁰	96
London	Observational	1972 – 1989 ⁴¹	169

* Comprises PSC cohorts ~ / ≥100 patients, which have contributed data to the international PSC Study Group (IPSCSG). Presented reports are likely to include those wherein more than one publication stems from a given cohort.

** Includes post-hoc outcome analysis of patients included in prior clinical trials.

*** Includes a subset of patients subject to an open-label study of endoscopic biliary intervention.

Supplementary References

- 1 Boberg KM, Rocca G, Egeland T, *et al.* Time-dependent Cox regression model is superior in prediction of prognosis in primary sclerosing cholangitis. *Hepatology* 2002; **35**: 652–7.
- 2 Boberg KM, Bergquist A, Mitchell S, *et al.* Cholangiocarcinoma in primary sclerosing cholangitis: risk factors and clinical presentation. *Scand J Gastroenterol* 2002; **37**: 1205–11.
- 3 Lindström L, Boberg KM, Wikman O, *et al.* High dose ursodeoxycholic acid in primary sclerosing cholangitis does not prevent colorectal neoplasia. *Aliment Pharmacol Ther* 2012; **35**: 451–7.
- 4 Lindström L, Hultcrantz R, Boberg KM, Friis-Liby I, Bergquist A. Association Between Reduced Levels of Alkaline Phosphatase and Survival Times of Patients With Primary Sclerosing Cholangitis. *Clin Gastroenterol Hepatol* 2013; **11**: 841–6.
- 5 Olsson R, Boberg KM, Schaffalitsky de Muckadell O, *et al.* High-Dose Ursodeoxycholic Acid in Primary Sclerosing Cholangitis: A 5-Year Multicenter, Randomized, Controlled Study. *Gastroenterology* 2005; **129**: 1464–72.
- 6 Vesterhus M, Hov JR, Holm A, *et al.* Enhanced liver fibrosis score predicts transplant-free survival in primary sclerosing cholangitis. *Hepatology* 2015; **62**: 188–97.
- 7 Benito de Valle M, Müller T, Björnsson E, *et al.* The impact of elevated serum IgG4 levels in patients with primary sclerosing cholangitis. *Dig Liver Dis* 2014; **46**: 903–8.
- 8 de Valle MB, Björnsson E, Lindkvist B. Mortality and cancer risk related to primary sclerosing cholangitis in a Swedish population-based cohort. *Liver Int* 2012; **32**: 441–8.




- 9 Zenouzi R, Weismüller TJ, Hübener P, *et al.* Low risk of hepatocellular carcinoma in patients with primary sclerosing cholangitis with cirrhosis. *Clin Gastroenterol Hepatol* 2014; **12**: 1733–8.
- 10 Tischendorf JJW, Hecker H, Krüger M, Manns MP, Meier PN. Characterization, Outcome, and Prognosis in 273 Patients with Primary Sclerosing Cholangitis: A Single Center Study. *Am J Gastroenterol* 2007; **102**: 107–14.
- 11 Zenouzi R, Weismüller TJ, Jørgensen KK, *et al.* No Evidence That Azathioprine Increases Risk of Cholangiocarcinoma in Patients With Primary Sclerosing Cholangitis. *Clin Gastroenterol Hepatol Off Clin Pract J Am Gastroenterol Assoc* 2016; **14**: 1806–12.
- 12 Jendrek ST, Gotthardt D, Nitzsche T, *et al.* Anti-GP2 IgA autoantibodies are associated with poor survival and cholangiocarcinoma in primary sclerosing cholangitis. *Gut* 2016; : gutjnl – 2016–311739.
- 13 Fevery J, Henckaerts L, Van Oirbeek R, *et al.* Malignancies and mortality in 200 patients with primary sclerosing cholangitis: a long-term single-centre study: Malignancies and mortality in PSC. *Liver Int* 2012; **32**: 214–22.
- 14 Fevery J, Van Steenberghe W, Van Pelt J, *et al.* Patients with large-duct primary sclerosing cholangitis and Crohn’s disease have a better outcome than those with ulcerative colitis, or without IBD. *Aliment Pharmacol Ther* 2016; **43**: 612–20.
- 15 Alswat K, Al-Harthy N, Mazrani W, Alshumrani G, Jhaveri K, Hirschfield GM. The spectrum of sclerosing cholangitis and the relevance of IgG4 elevations in routine practice. *Am J Gastroenterol* 2012; **107**: 56–63.

- 16 Garioud A, Seksik P, Chrétien Y, *et al.* Characteristics and clinical course of primary sclerosing cholangitis in France: a prospective cohort study. *Eur J Gastroenterol Hepatol* 2010; **22**: 842–7.
- 17 Gotthardt DN, Rudolph G, Klöters-Plachky P, Kulaksiz H, Stiehl A. Endoscopic dilation of dominant stenoses in primary sclerosing cholangitis: outcome after long-term treatment. *Gastrointest Endosc* 2010; **71**: 527–34.
- 18 Rudolph G, Gotthardt D, Kloeters-Plachky P, Rost D, Kulaksiz H, Stiehl A. In PSC with dominant bile duct stenosis, IBD is associated with an increase of carcinomas and reduced survival. *J Hepatol* 2010; **53**: 313–7.
- 19 Rupp C, Rössler A, Halibasic E, *et al.* Reduction in alkaline phosphatase is associated with longer survival in primary sclerosing cholangitis, independent of dominant stenosis. *Aliment Pharmacol Ther* 2014; **40**: 1292–301.
- 20 Rupp C, Bode KA, Chahoud F, *et al.* Risk factors and outcome in patients with primary sclerosing cholangitis with persistent biliary candidiasis. *BMC Infect Dis* 2014; **14**: 562.
- 21 Gauss A, Sauer P, Stiehl A, *et al.* Evaluation of Biliary Calprotectin as a Biomarker in Primary Sclerosing Cholangitis. *Medicine (Baltimore)* 2016; **95**. DOI:10.1097/MD.00000000000003510.
- 22 Okolicsanyi L, Fabris L, Viaggi S, Carulli N, Podda M, Ricci G. Primary sclerosing cholangitis: clinical presentation, natural history and prognostic variables: an Italian multicentre study. The Italian PSC Study Group. *Eur J Gastroenterol Hepatol* 1996; **8**: 685–91.

- 23 Boonstra K, Weersma RK, van Erpecum KJ, *et al.* Population-based epidemiology, malignancy risk, and outcome of primary sclerosing cholangitis. *Hepatology* 2013; **58**: 2045–55.
- 24 Claessen MMH, Vleggaar FP, Tytgat KMAJ, Siersema PD, van Buuren HR. High lifetime risk of cancer in primary sclerosing cholangitis. *J Hepatol* 2009; **50**: 158–64.
- 25 Ponsioen CY, Vrouenraets SME, Prawirodirdjo W, *et al.* Natural history of primary sclerosing cholangitis and prognostic value of cholangiography in a Dutch population. *Gut* 2002; **51**: 562–6.
- 26 Ponsioen CY, Lam K, van Milligen de Wit AW, Huibregtse K, Tytgat GN. Four years experience with short term stenting in primary sclerosing cholangitis. *Am J Gastroenterol* 1999; **94**: 2403–7.
- 27 Ponsioen CY, Reitsma JB, Boberg KM, Aabakken L, Rauws EA, Schrumpf E. Validation of a cholangiographic prognostic model in primary sclerosing cholangitis. *Endoscopy* 2010; **42**: 742–7.
- 28 Broomé U, Olsson R, Lööf L, *et al.* Natural history and prognostic factors in 305 Swedish patients with primary sclerosing cholangitis. *Gut* 1996; **38**: 610–5.
- 29 Bergquist A, Said K, Broomé U. Changes over a 20-year period in the clinical presentation of primary sclerosing cholangitis in Sweden. *Scand J Gastroenterol* 2007; **42**: 88–93.
- 30 Bergquist A, Ekbom A, Olsson R, *et al.* Hepatic and extrahepatic malignancies in primary sclerosing cholangitis. *J Hepatol* 2002; **36**: 321–7.


- 31 Bergquist A, Glaumann H, Persson B, Broomé U. Risk factors and clinical presentation of hepatobiliary carcinoma in patients with primary sclerosing cholangitis: A case-control study. *Hepatology* 1998; **27**: 311–6.
- 32 Imam MH, Silveira MG, Sinakos E, *et al.* Long-term Outcomes of Patients With Primary Biliary Cirrhosis and Hepatocellular Carcinoma. *Clin Gastroenterol Hepatol* 2012; **10**: 182–5.
- 33 Lindor KD, Kowdley KV, Luketic VAC, *et al.* High-dose ursodeoxycholic acid for the treatment of primary sclerosing cholangitis. *Hepatology* 2009; **50**: 808–14.
- 34 Eaton JE, Silveira MG, Pardi DS, *et al.* High-dose ursodeoxycholic acid is associated with the development of colorectal neoplasia in patients with ulcerative colitis and primary sclerosing cholangitis. *Am J Gastroenterol* 2011; **106**: 1638–45.
- 35 Imam MH, Thackeray EW, Lindor KD. Colonic neoplasia in young patients with inflammatory bowel disease and primary sclerosing cholangitis. *Colorectal Dis* 2013; **15**: 198–203.
- 36 Wiesner RH, Grambsch PM, Dickson ER, *et al.* Primary sclerosing cholangitis: natural history, prognostic factors and survival analysis. *Hepatology* 1989; **10**: 430–6.
- 37 Burak K, Angulo P, Pasha TM, Egan K, Petz J, Lindor KD. Incidence and Risk Factors for Cholangiocarcinoma in Primary Sclerosing Cholangitis. *Am J Gastroenterol* 2004; **99**: 523–6.

- 38 Toy E, Balasubramanian S, Selmi C, Li C-S, Bowlus CL. The prevalence, incidence and natural history of primary sclerosing cholangitis in an ethnically diverse population. *BMC Gastroenterol* 2011; **11**: 83.
- 39 Chapman MH, Webster GJ, Bannoo S, Johnson G, Wittmann J, Pereira SP. Cholangiocarcinoma and dominant strictures in patients with primary sclerosing cholangitis; a 25 year single centre experience. *Eur J Gastroenterol Hepatol* 2012; **24**: 1051–8.
- 40 Marelli L, Xirouchakis E, Kalambokis G, Cholongitas E, Hamilton MI, Burroughs AK. Does the severity of primary sclerosing cholangitis influence the clinical course of associated ulcerative colitis? *Gut* 2011; **60**: 1224–8.
- 41 Farrant JM, Hayllar KM, Wilkinson ML, *et al.* Natural history and prognostic variables in primary sclerosing cholangitis. *Gastroenterology* 1991; **100**: 1710–7.

A **Cholangiocarcinoma - overall****B** **Cholangiocarcinoma - after the 1st year****C** **Cholangiocarcinoma - within the 1st year**

Supplementary Figure 2


A



—	254	165	84	30	10	Pts. at risk
—	470	261	116	51	21	Pts. at risk
—	6397	3756	1930	861	342	Pts. at risk

[Click here to download Supporting Document Supplementary Figure 2 boxed.pdf](#)

B



—	247	161	81	29	9	Pts. at risk
—	460	254	111	46	17	Pts. at risk
—	6122	3542	1802	789	308	Pts. at risk